

Los Alamos archives: gr-qc/9411051
 © H.C. Rosu, Nuovo Cimento B 108 (1993) 313-329

HYBRIDIZING THE SKYRMION WITH AN ANTI-DE SITTER BAG

HARET C. ROSU

Instituto de Física, Universidad de Guanajuato, Apdo Postal E-143, León, Gto, Mexico
rosu@ifug3.ugto.mx

I discuss a phenomenological model of the nucleon in which a small anti-de Sitter bag is placed into the Skyrmion configuration. Such a bag has a timelike boundary and allows naturally the Cheshire Cat Principle. Very important in this model is the membrane of the bag, the 3-dimensional time-space manifold $S^1 \times S^2$, in which topological techniques will come into play.

Contents

1. Introduction
2. Review of de Sitter and anti-de Sitter spaces
3. AdS femtocosmology
4. Cheshire Cat Principle and the AdS bag
5. The pion and the AdS bag
6. Conclusions and perspectives

References

This is a slightly revised version of IC-26/92 and of Nuovo Cimento B 108, 313-329 (1993). IC-26/92 has been written in January 1992, and revised in October 1992 and November 1994. The companion paper is Nuovo Cimento A 107, 1543-1547 (1994) [gr-qc/9406017(IFUG-17b/94)].

IFUG-17a/94 [$\mathcal{HCR}$]

1. Introduction

Since 1932 the atomic nuclei are considered to be made of protons and neutrons. This is due mainly to the fact that their degree of virtuality is very low (8 MeV as compared to 939 MeV). Nevertheless quantum chromodynamics (QCD) ¹, arising so powerfully in the seventies to explain hadronic structure, forced us to look at hadrons and atomic nuclei as merely low energy phenomenology of that theory.

A decisive step forward on such a route was the MIT bag model, which subsequently gave birth to an impressive number of other bags. Most remarkably, a model of the fifties due to Skyrme became prominent recently ². As it is well known the attractive features of the Skyrmion are connected with some fresh mathematics streaming into the realm of hadron and nuclear physics from such fields like topology and differential geometry. At the same time, Skyrmions have been proven to have a good matching to Yukawa theory as well as to bag models through hybridization. Thus, Skyrmions may be looked upon as a necessary development in low energy hadronic and nuclear phenomenology. In the past a number of important results were established within Skyrmion physics, some of which are :

- i) Skyrme model is an effective low energy/large N_c limit of QCD ³.
- ii) The baryon number has a topological nature and belongs to the homotopy group $\pi_3(S^3) = \mathbb{Z}$ ⁴. The winding baryon number of the mappings $S^3 \rightarrow S^3$ is

$$B = \int d^3x B_0 , \quad (1.1)$$

where B_0 is the time component of the conserved topological current

$$B_\mu = \frac{1}{24\pi^2} \epsilon_{\mu\nu\alpha\beta} \text{tr} L^\nu L^\alpha L^\beta . \quad (1.2)$$

The L matrices, $L_\mu = U^\dagger \partial_\mu U$, are left topological currents of $SU(2)$ matrices parametrizing the pion fields

$$U(x) = \exp[2i\tau^a \phi_a / F_\pi] , \quad (1.3)$$

where τ^a are the Pauli matrices, ϕ_a are the pion fields, and F_π is the pion decay constant used to normalize the pion fields.

- iii) The Wess-Zumino term added to the Skyrme Lagrangian fuses two unwanted discrete symmetries making the model viable ⁵.

The Skyrme Lagrangian is invariant separately under

$$P_1 : \vec{x} \rightarrow -\vec{x} \quad (1.4)$$

and

$$P_2 : U \rightarrow U^\dagger , \quad (1.5)$$

which are combined into the parity transformation $P = P_1 \cdot P_2$. However, processes like $K^+ K^- \rightarrow \pi^+ \pi^- \pi^0$ break P_1 and P_2 . Therefore an action term is required,

which breaks these symmetries keeping the combined parity P invariant. The Wess-Zumino-Witten (WZW) term is acting like that for the SU(3) Skyrmions. It is a 5-dimensional integral over a disk D whose boundary is the four dimensional space-time

$$S_{WZW} = -\frac{iN_c}{240\pi^2} \int_D d^5x \epsilon^{\alpha\beta\gamma\delta\epsilon} \text{Tr} L_\alpha L_\beta L_\gamma L_\delta L_\epsilon . \quad (1.6)$$

The coefficient in front of the integral, proportional to N_c , is determined by computing electromagnetic processes such as $\pi^0 \rightarrow 2\gamma$, for this action, with an additional gauge-invariant coupling to the electromagnetic field, and comparing it with the direct QCD calculation ⁶.

iv) The Cheshire Cat Principle: if a bag is introduced into the Skyrme soliton as a kind of defect ⁷, then the dynamics is independent of the boundary conditions at the bag surface. This is an exact statement in two dimensions only ⁸.

Usually, one will make use of a chiral coupling between quarks and pions at the bag surface, i.e., $U_5 = \exp(i\vec{\tau} \cdot \vec{\phi}\gamma_5/f_\pi)$.

In the following we want to introduce into the Skyrmon a rather special bag possessing anti-de Sitter (AdS) geometry and to make an analysis of the consequences of proceeding in this way. The first two sections of the paper are short surveys on the topics of de Sitter (dS) spaces and strong gravity (femtocosmology). They were written for the convenience of the readers (who are supposed to belong mostly to the Skyrmon society). The third section shows that the AdS bag is very appealing for the Cheshire Cat Principle. In the fourth section we comment on the pion as the AdS Goldstone boson. Finally, we add some conclusions. The present paper is an updated and improved version of a talk delivered at Poiana Brasov School in the summer of 1988 ⁹.

2. Review of de Sitter and anti-de Sitter spaces

de Sitter spaces are “respectable” objects in theoretical physics ever since 1917. There is an extensive literature on various aspects concerning these spacetimes. Very good introductory material could be found in a number of well known books and review papers ¹⁰.

de Sitter universes are maximally symmetric solutions of Einstein cosmological equations

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} + \Lambda g_{\mu\nu} = \kappa T_{\mu\nu} \quad (2.1) ,$$

with no energy-momentum tensor, $T_{\mu\nu} = 0$. Thus, de Sitter solutions are vacuum ones, and it was precisely this feature as related to a variant of Mach principle which was very much discussed in 1917, when W. de Sitter introduced his universe in Physics ¹¹.

On the other hand, spaces of constant negative curvature were brought into Physics by A.A. Friedman in his second famous cosmological paper of 1924 written at the suggestion of his good friend Tamarkin. Mathematically, these spaces had already at that time a history of one hundred years ¹², starting with Lobachevsky and

Bolyai, although Minding was the first one who studied surfaces of constant negative curvature (1839), and of course we must remember Beltrami who established the connection between hyperbolic geometry and these surfaces in 1868.

de Sitter space was much investigated in cosmology, as it was the first cosmological model possessing a red-shift (i.e., a direct observable). After 1980, many authors have dealt with this space in connection with the inflationary hypothesis asserting that a dS phase dominated the very early Universe ($t \leq 10^{-35}$ s; $T \geq 10^{15}$ GeV).

AdS space, on the other hand, turned out to be very important in extended supergravity, in Kaluza-Klein theories, and more generally in microcosmos. This is due to a number of reasons, amongst which positive definiteness of the energy operator, the causal relationship one could set up in the AdS case, the possibility to construct an S-matrix and also the analogue of Penrose transform to a generalized twistor space.

Considered as Riemannian spaces, dS universes being of constant curvature have the curvature tensor given by the following formula

$$R_{\mu\nu\rho\sigma} = a^2(g_{\mu\rho}g_{\nu\sigma} - g_{\mu\sigma}g_{\nu\rho}) . \quad (2.2)$$

Locally, dS spaces are expressed by the equation

$$R_{\mu\nu} - 1/4 R g_{\mu\nu} = 0 \quad (2.3)$$

In mathematics such spaces are called Einstein spaces¹³. If, as physicists, we prefer to keep a cosmological constant in our line of reasoning, then we have the following well-known relation

$$\Lambda = 1/4R . \quad (2.4)$$

Assuming spherical symmetry, de Sitter line element turns into the form first written by de Sitter as follows

$$ds^2 = (1 - 1/3\Lambda r^2)dt^2 - (1 - 1/3\Lambda r^2)^{-1}dr^2 - r^2(d\theta^2 + \sin^2\theta d\phi^2) . \quad (2.5)$$

The embedding in a 5-dimensional pseudoeuclidean space is given in the case $\Lambda > 0$ by

$$\begin{aligned} x_0 &= \left(\frac{3}{\Lambda}\right)^{\frac{1}{2}} \left(1 - \frac{1}{3}\Lambda r^2\right)^{\frac{1}{2}} \sinh\left[\left(\frac{\Lambda}{3}\right)^{\frac{1}{2}} t\right] \\ x_1 &= \left(\frac{3}{\Lambda}\right)^{\frac{1}{2}} \left(1 - \frac{1}{3}\Lambda r^2\right)^{\frac{1}{2}} \cosh\left[\left(\frac{\Lambda}{3}\right)^{\frac{1}{2}} t\right] \\ x_2 &= r \sin\theta \cos\phi \\ x_3 &= r \sin\theta \sin\phi \\ x_4 &= r \cos\theta . \end{aligned} \quad (2.6)$$

These embedding equations turn the dS universe into the hypersurface

$$+x_0^2 - x_1^2 - x_2^2 - x_3^2 - x_4^2 = -\frac{3}{\Lambda} . \quad (2.7)$$

For the AdS space ($\Lambda < 0$) the embedding formulas are

$$\begin{aligned}
x_0 &= \left(\frac{3}{-\Lambda}\right)^{\frac{1}{2}} \left(1 - \frac{1}{3}\Lambda r^2\right)^{\frac{1}{2}} \sin\left[\left(\frac{-\Lambda}{3}\right)^{\frac{1}{2}} t\right] \\
x_1 &= \left(\frac{3}{-\Lambda}\right)^{\frac{1}{2}} \left(1 - \frac{1}{3}\Lambda r^2\right)^{\frac{1}{2}} \cos\left[\left(\frac{-\Lambda}{3}\right)^{\frac{1}{2}} t\right] \\
x_2 &= r \sinh \theta \cos \phi \\
x_3 &= r \sinh \theta \sin \phi \\
x_4 &= r \cosh \theta .
\end{aligned} \tag{2.8}$$

The pseudoeuclidean coordinates are constrained this time to the hypersurface

$$+x_0^2 - x_1^2 - x_2^2 - x_3^2 + x_4^2 = -\frac{3}{\Lambda} . \tag{2.9}$$

The four-dimensional quadrics (2.7) and (2.9) are a four-hyperboloid with one sheet in the dS case and a four-hyperboloid with two sheets in the AdS case. Their topology is $R \times S^3$ and $S \times R^3$, respectively.

The line element (2.5) could also be written in the Robertson-Walker (expansion) form

$$ds^2 = d\tau^2 - \frac{3}{\Lambda} \cosh^2\left[\left(\frac{\Lambda}{3}\right)^{\frac{1}{2}} \tau\right] (d\chi^2 + \sin^2 \chi d\Omega^2), \text{ for } \Lambda > 0 \tag{2.10}$$

$$ds^2 = d\tau^2 + \frac{3}{\Lambda} \cos^2\left[\left(\frac{-\Lambda}{3}\right)^{\frac{1}{2}} \tau\right] (d\chi^2 + \sinh^2 \chi d\Omega^2), \text{ for } \Lambda < 0 . \tag{2.11}$$

The above expressions make it obvious that dS universe is closed, i.e., with a spherical 3-dimensional geometry, while the AdS universe is open, i.e., with a hyperbolic 3-dimensional geometry.

For the AdS case, it is worth noting that the scale factor (the radius of the universe in the older literature) is an oscillating function, with a maximum amplitude

$$r_{max} = \left(\frac{3}{-\Lambda}\right)^{\frac{1}{2}} . \tag{2.12}$$

The last useful form of the line element is the conformal one

$$ds^2 = (a \cos \rho)^{-2} [dt^2 - d\rho^2 - \sin^2 \rho (d\theta^2 + \sin^2 \theta d\phi^2)] , \tag{2.13}$$

where the radial variable has the range $0 < \rho < \pi/2$, the other ranges being the common ones. The parameter a is given by $R = -12a^2$.

As usual when looking to spacetimes as 4-dimensional geometries, we have to consider first of all their kinematical group of isometries. The de Sitter and Anti-de Sitter groups are the most simple and direct generalization of the Poincare group, which can be obtained from the first ones by Inönü-Wigner contraction procedure.

In homogeneous coordinates, the Absolute (i.e., the invariant quadric) of the Lorentz group is given by

$$+\xi_1^2 + \xi_2^2 + \xi_3^2 - c^2 \xi_4^2 = 0 , \tag{2.14}$$

whereas the Absolutes of dS/AdS groups are determined by

$$+\xi_1^2 + \xi_2^2 + \xi_3^2 - c^2 \xi_4^2 \pm R^2 \xi_5^2 = 0 . \quad (2.15)$$

One can see immediately that this generalization of the Lorentz group is equivalent to the minimal introduction of cosmology through a constant curvature of positive value for dS space and of negative value for AdS space.

We shall use mostly geometrical units: $c=1; \Lambda = 3; \hbar = 1$.

The quadratic form (2.9) corresponding to the AdS space has to be invariant under the action of the isometry group $SO(3,2)$. The ten Killing vectors are of the usual “angular momentum” type

$$K_{AB} = -(x_A \partial_B - x_B \partial_A) \quad (2.16)$$

and satisfy the Lie algebra

$$[K_{AB}, K_{CD}] = +K_{AC}\eta_{BD} + K_{BC}\eta_{AD} - K_{AD}\eta_{BC} - K_{BD}\eta_{AC} , \quad (2.17)$$

where $\eta_{AB} = \text{diag}(+1, -1, -1, -1, +1)$. When one is passing to the quantum theory the covariant generators have to be used

$$\hat{M}_{AB} = -iK_{AB} . \quad (2.18)$$

To be more specific one has noncommutative translations

$$[\hat{P}_\alpha, \hat{P}_\beta] = R\hat{M}_{AB} . \quad (2.19)$$

If we denote

$$\hat{M}_{A4} = r\hat{P}_a , \quad (2.20)$$

where r is the AdS radius of curvature, and

$$\hat{M}_{AB} = \hat{M}_{\alpha\beta} \quad (2.21)$$

for the other generators, we get the compact Lie product (commutator) of the type (2.17). Inönü-Wigner contraction simply means $r \rightarrow 0$.

Since $SO(3,2)$ is a noncompact group, the hermiticity of the operators, ($\hat{M}_{AB} = \hat{M}_{AB}^\dagger$), implies the unitarity of only the infinite dimensional representations. The compact operators are $\hat{M}_{04}$ and $\hat{M}_{ab}$ and the noncompact ones are $\hat{M}_{0a}$ and $\hat{M}_{a4}$. The corresponding physical quantities are obtained from the contracted Lie algebra (the Poincaré one). Since under contraction $\hat{M}_{04}$ goes into $\hat{P}_0$, which is the time translation operator, $\hat{M}_{04}$ could be identified with the energy operator; also $\hat{M}_{ab}$ are angular momentum operators; the noncompact operators $\hat{M}_{0a}$ and $\hat{M}_{a4}$ correspond to the momentum operators and Lorentz boost operators, respectively. The center of $SO(3,2)$ is bidimensional. The two Casimir operators are the following

$$\hat{C}_2 = 1/2 \hat{M}^{AB} \hat{M}_{AB}$$

$$\hat{C}_4 = 1/2 \hat{M}^{AB} \hat{M}_{BC} \hat{M}^{CD} \hat{M}_{DE} . \quad (2.22)$$

The Casimir operators of $SO(3,2)$ are similar to the Casimir operators of the Poincaré group. In the representations of the Poincaré group, the latter ones have the eigenvalues m^2 and $m^2 s(s+1)$, where m denotes the mass and s the spin. However the $SO(3,2)$ representations are not usually labeled by means of $\hat{C}_2$ and $\hat{C}_4$ eigenvalues. Instead of this, E_0 , the smallest value of the AdS energy (in units of a) and s , the minimal eigenvalue of the total angular momentum are used (they correspond to the maximal compact subgroup of $SO(3,2)$ which is $SO(3,2) \otimes SO(3,2)$). The reason for that is to be found in the fact that the Casimir eigenvalues do not uniquely specify the representations of noncompact groups. Another general fact for such groups is the occurrence of indecomposable representations (noncompletely reducible), of much help in doing generalized dS quantum electrodynamics in parallel to the usual Minkowski QED ¹⁴.

The eigenvalue of the Casimir operator $\hat{C}_2$ could be expressed in terms of E_0 and s as follows

$$C_2 = E_0(E_0 - 3) + s(s+1) . \quad (2.23)$$

Having two timelike directions the group $O(3,2)$ must contain as a factor group the conformal group of 3 dimensions, i.e., the group of symmetries of a cone in a 3-dimensional pseudo-Euclidian space with signature $(+, -, -)$. Consider the 5-cone ($\eta_{AB} x^A x^B = 0$) and define homogeneous coordinates in the following way

$$y^\mu = x^\mu / (x_3 + x_4) , \quad (2.24)$$

where $\mu = 0, 1, 2$. The linear action of $O(3,2)$ on the 5-cone given by

$$x'_A = L^B_A x_B , \quad (2.25)$$

with $L^B_A \in O(3,2)$ induces by (2.24) a nonlinear action on the Minkowski space of y_μ denoted as $\text{Mink}(y)$. The action of $O(3,2)$ on $\text{Mink}(y)$ is not effective, i.e., we have two identity transformations on $\text{Mink}(y)$ coming from $O(3,2)$, namely the subgroup $Z_2 = \{\Lambda \in O(3,2) : \Lambda x = \pm x\}$. The conformal group is defined as: $C(1,2) \sim O(3,2)/Z_2$. The group $O(3,2)$ has four components. That one connected to the identity is

$$SO_0(3,2) = \{L \in O(3,2) : \det L = 1; L^0_0 L^4_4 - L^4_0 L^0_4 > 1\} . \quad (2.26)$$

The other components are obtained through combined reversals of the signs of $\det L$ and of the “time order”. One can see that $SO(3,2) \subset C(1,2)$. To pass from the component connected to the identity to some other component, discrete operations are required

$$\mathcal{T} := \{x^0 \rightarrow -x^0; x^4 \rightarrow x^4; x^i \rightarrow x^i\} \quad (2.27)$$

$$\mathcal{P} := \{x^0 \rightarrow x^0; x^4 \rightarrow x^4; x^i \rightarrow -x^i\} \quad (2.28)$$

$$\mathcal{A} := \{x^0 \rightarrow -x^0; x^4 \rightarrow -x^4; x^i \rightarrow -x^i\} . \quad (2.29)$$

$\mathcal{T}, \mathcal{P}$, and $\mathcal{A}$ stand for time reversal, space parity, and antipodal inversion (or parity), respectively. $\mathcal{A}$ commutes with all elements of $SO(3,2)$ and has the involutory eigenvalues ± 1 . $\mathcal{P}$ and $\mathcal{A}$ are related by

$$\mathcal{P} = \mathcal{V} \mathcal{A} \mathcal{V}^{-1} , \quad (2.30)$$

where $\mathcal{V} \in SO_0(3,2)$ is given by

$$\mathcal{V} := \{x^0 \rightarrow -x^0; x^4 \rightarrow -x^4; x^i \rightarrow x^i\} . \quad (2.31)$$

For the spin-1/2 fields in dS space we have to add a “spin part” to the $SO(3,2)$ “angular part”

$$\hat{J}_{AB} = \hat{M}_{AB} + \hat{S}_{AB} . \quad (2.32)$$

The charge conjugation matrix of AdS space is the standard 4×4 matrix of Minkowski space ¹⁵. For more sophisticated applications in supergravity and KK theory where AdS manifolds occur as natural backgrounds, it is worth noting that the SUSY extension of $SO(3,2)$, which is the supergroup $Osp(4/N)$ has $4N$ Killing spinors playing the same role in the spinor space as do Killing vectors in the boson space. In the case $N=1$, all linear unitary irreducible representations of AdS SUSY have been classified by Heidenreich ¹⁶. The best (super)review has been written in 1986 ¹⁷. The most complete discussion of the $SO_0(3,2)$ representations has been provided by Angelopoulos ¹⁸. For more physics-oriented treatments one should see Gibbons ¹⁵, as well as Nicolai ¹⁹.

The representations $D(E_0, s)$ are unitary iff

$$\begin{aligned} E_0 &\geq s + 1, \quad s = 1, 3/2, 2, 5/2, \dots \\ E_0 &\geq s + 1/2, \quad s = 0, 1/2 . \end{aligned} \quad (2.33)$$

Fields with $s \geq 1$ reach the unitarity limit when $E_0 = s+1$ and since they contract to the massless discrete helicity representations of the Poincaré group, they are called “massless”. Fields with $s=0, 1/2$ have $E_0 = s + 1/2$ in the unitary limit but have no Poincaré limit. They are not massless and have been called singletons by Dirac ²⁰, who discovered them. As a tribute, Fronsdal ²¹, dubbed the boson singleton as the RAC (i.e., $D(1/2,0)$), and the fermion one as the DI, (i.e., $D(1,1/2)$). They have been under the focus of many authors, especially in their superform since they could be related to super p-branes ²². From the group-theoretical standpoint (super)singletons are included in an ultrashort (super)multiplet.

Singletons have two remarkable properties namely, the one-singleton states are unobservable and the two-singleton states are always massless, i.e.,

$$Singleton \otimes Singleton = \oplus Massless Reps . \quad (2.34)$$

The essential point to stress is that the quantum field theory of singletons is intimately related to the spatial infinity of AdS space which is the famous manifold $S^1 \times S^2$. Geometrically it is precisely this AdS boundary that give birth to singleton

fields. The fact that the AdS boundary is the remarkable 3-dimensional $S^1 \times S^2$ manifold makes us foresee many topological methods entering the field. The very interesting and well-developed topology (in Mathematics) of this “Membrane at the End of the Universe”²³, is at the same time the topology of the “Nucleon Membrane”, as I would call it.

We have to recall also the strong analogy between quarks and singletons which has been put forth by Flato and Fronsdal²⁴, and as a matter of fact, one can look at them as completely equivalent objects. This is, in our opinion, the most important property of AdS space still to be fully exploited²⁵.

Many encumbrances reveal themselves when someone is going to tackle the problem of quantizing fields in AdS background. This is due to the $S^1 \times R^3$ topology of AdS. Avis, Isham, and Storey²⁶ have discussed in a very clear way the quantization of scalar fields in AdS space. The things to be emphasized are first, the impossibility to specify Cauchy data on a single spatial hypersurface, in other words, AdS is a non-global hyperbolic manifold. Second, it possesses closed timelike curves. This is considered as an unwanted property and is avoided by going to AdS universal covering space (CAdS) which is a global manifold with a $R \times S^3$ topology allowing to be conformally mapped onto half of the Einstein static universe (ESU). The boundary at spatial infinity remains timelike even for the CAdS case implying loss or gain of information through spatial infinity in finite coordinate time. Such a situation is unacceptable for a well defined quantum field theory. In order to develop a consistent quantization scheme Avis, Isham, and Storey have introduced reflective boundary conditions (RBC) at spatial infinity, in other words, they have put there reflecting walls. We shall use again results from²⁶ in section 3.

Until about five years ago, AdS quantum field theory has been a flourishing topic, especially due to SUSY, SUGRA, and KK theories²⁷, but meanwhile the activity in the field has diminished as the result of other fashionable discoveries. We recall here the parallel and less ambitious use of dS groups and their SUSY extensions as dynamical (spectrum generating) groups²⁸. In this case one is not interested so much in the structure of the “constituents”, but rather in their relative motions and for that purpose the oscillator representations are recommendable. AdS space may be considered as a general relativistic version of the harmonic oscillator potential with the curvature playing the role of the harmonic oscillator coupling constant. In this way, dynamical spectra (e.g., Regge trajectories) are reproducible by means of AdS dynamical group. On the other hand, dS groups have also been used as gauge groups²⁹, and more practically in some powerful techniques for separation of variables in partial differential equations³⁰.

3. AdS femtocosmology

As has been put by Allen and Turyn, “de Sitter spaces interest different people for different reasons”³¹. Our goal is to make a connection between AdS physics and Skyrmion physics. This could be done by inserting into the Skyrmion configuration a bag (or a defect in the terminology of G. Brown) possessing AdS geometry. It is

a kind of (femto)cosmology, which is known to have some tradition ³².

An AdS bag was beautifully discussed for the first time by Salam and Strathdee in the f-g theory context ³³. Among precursors, we name Prasad, who developed somewhat similar ideas before the Bag Era ³⁴. Subsequently, a dS bag was compactly described by Haba ³⁵. Romaniuk studied an AdS bag with “dust” as the gravitational source ³⁶. Important developments on the AdS geometrical confinement problem have been worked out by the Nijmegen group ³⁷.

Femtocosmology could be considered also as a variant of the bimetric general relativity ³⁸, applied at the hadron spacetime scale where the coexistence of the two metrics is supposed to occur, the common Minkowski one and the AdS one.

The basic idea in femtocosmology of considering hadrons as strongly curved femtouniverses is accomplished by scaling down the AdS Universe by giving to the curvature scalar a value fixed rather arbitrarily by some hadronic scale, or equivalently by introducing a huge cosmological constant

$$\Lambda_h \sim 10^{26} \text{cm}^{-2} \sim 10^{-2} \text{GeV}^2 . \quad (3.1)$$

It is worth noting that such a huge gravitational effect of Higgs origin is nevertheless considered small on the scale of quantum gravity. We are still at several orders of magnitude away from the Planck scale.

What is the reason making possible the scaling up and/or down of universes? In other terms, what does make the curvature scalar arbitrary? The answer to these important questions is given by the hyperbolic geometry. Let us take the hyperbolic surface $\mathcal{H}^2$, though the discussion holds good for any number of dimensions. It is a mathematical fact that the angle of parallelism in $\mathcal{H}^2$ is given by the well-known relation

$$\tan[1/2\Pi(x)] = e^{-x/\kappa} , \quad (3.2)$$

where κ is the hyperbolic curvature, which one could express more intuitively in terms of the distance $\mathcal{D}$ corresponding to an angle of parallelism of $\pi/4$ radians. The connection is

$$\kappa = 0.2\mathcal{D} . \quad (3.3)$$

The point is that hyperbolic geometry asserts the completely arbitrary character of the distance $\mathcal{D}$. It follows that Mathematics allows the scaling of universes of constant curvature. In AdS spaces the radius of curvature appears on the same level as the velocity of light, as one can see from (2.15). Hence it may be considered as a constant of physics very similar to a fundamental length ³⁹. There is no etalon for the curvature, and consequently it is difficult to consider it as a physical constant. Its nature is purely geometrical. However the spontaneous breaking process is the common procedure by which geometrical constants are introduced into physics. Since AdS group is a subgroup of the conformal group $C(4,2)$, the spontaneous breaking of the conformal symmetry leaves only $SO(3,2)$ symmetry in the real world. This symmetry breaking was studied in a very straightforward way by Fubini ⁴⁰. On the other hand, de Sitter spaces are in turn conformally

flat and therefore a rescaled conformally coupled field does not recognize as special the length scale set by the curvature of the spacetime in which it lives up to the conformal anomaly.

In order to fix the existing value $r \sim 1$ fm, we shall proceed rather arbitrarily and use a grand-unified Higgs field. Indeed it is essential to note that one can get the AdS geometry by means of a scalar field with a huge mass, since in this case the scalar energy-momentum tensor simulates the vacuum one, more exactly in the limit $(\dot{\varphi}/\varphi)^2 \ll M^2$ we get the vacuum equation of state $p = -\epsilon$.

The following procedure is well known in relation to the spontaneous symmetry breaking rules ⁴¹. The Lagrangian of a complex scalar field is written down

$$\mathcal{L} = \partial_\mu \varphi^+ \partial^\mu \varphi + \mu^2 \varphi^+ \varphi - \lambda (\varphi^+ \varphi)^2 \quad (3.4)$$

The field $\varphi = \varphi_1 + i\varphi_2$ has hermitic components. One of them has a nonzero vacuum expectation value denoted by σ . Then the fields are shifted by σ . The new Lagrangian in the shifted fields ϕ_1 and ϕ_2 will be

$$L = \hat{L} + 1/2 \mu^2 \sigma^2 - 1/4 \lambda \sigma^4 . \quad (3.5)$$

All the operator terms have been included in $\hat{L}$. Stability arguments give

$$\sigma^2 = \mu^2 / \lambda . \quad (3.6)$$

The vacuum expectation value of the Hamiltonian will be

$$\langle H \rangle_{vac} = -\mu^2 \sigma^2 / 4 = \langle T_{\mu\nu} \rangle_{vac} . \quad (3.7)$$

Now, one could use Einstein equations to get

$$\langle T_{\mu\nu} \rangle_{vac} = \Lambda / \kappa . \quad (3.8)$$

From (3.4) we can get a negative cosmological constant of the form

$$\Lambda = -1/4 \mu^2 \sigma^2 . \quad (3.9)$$

The maximal value of the AdS radius will follow as

$$\mathcal{R} = (12/\mu^2 \sigma^2)^{1/2} . \quad (3.10)$$

In order to have an AdS vacuum at the hadron scale we need a massive GUT Higgs field with $\mu \sim 10^{15} GeV$. The Higgs pressure is given by the quantity Λ/κ .

4. Cheshire Cat Principle and the AdS bag

Throughout this section we shall stay in two dimensions. Cheshire Cat Principle (CCP) is a rigorous result in this case and has to do with the boundary conditions at the bag surface. CCP has still to be generalized to four dimensions. We recall that quantum field theory in two dimensions is dominated by the remarkable complete

equivalence of fermion and boson theories⁴². CCP which has been put forth in 1985 is just a consequence of this general feature⁸. For fermion-boson equivalence in four dimensions see the appendix in the paper of Rubakov⁴³.

Our physical scenario is that of a skyrmion with an AdS bag inside it. The 2-dimensional AdS space (AdS₂) has the following metric in intrinsic coordinates t, ρ

$$ds^2 = (a \cos \rho)^{-2} [dt^2 - d\rho^2] , \quad (4.1)$$

where $-\pi/2 < \rho < \pi/2$ and $-\pi \leq t \leq \pi$ in AdS, the latter range extending to the whole real axis for CAdS. The image of AdS₂ in Minkowski space is a finite spatial interval whose boundary consists of 2 points, $\rho = \pm\pi/2$.

The CCP concept has been introduced and developed at Nordita⁸; see also⁴⁴. We shall discuss only the abelian case, i.e., fermions of only one colour and flavor. CCP is opposite to confinement allowing fermions to leak out of the bag (in the membrane region in our model) and finally in the skyrmion region they show up as a soliton of the boson representation. It is some highly nontrivial topological dynamics in the $S^1 \times S^2$ membrane which is involved in the CCP process. The idea of the standard CCP is to make the Euler-Lagrange equations of the total action to coincide at least partially with the bosonization relations. This is achieved if one adds to the fermionic and bosonic action a boundary term of the form⁴⁵

$$S_B = \int_B d\epsilon^\mu (1/2 n_\mu \bar{\psi} e^{i\sqrt{4\pi}\gamma_5 \phi} \psi) , \quad (4.2)$$

where $d\epsilon^\mu = d\epsilon \cdot n^\mu$ is a surface area element, n^μ being the bag normal. Setting the total surface variation equal to zero, one can get the following two boundary equations

$$\begin{aligned} i \not{n} \psi &= n^2 e^{i\sqrt{4\pi}\gamma_5 \phi} \phi \psi \\ n \partial \phi &= \sqrt{\pi} \bar{\psi} \not{n} \gamma_5 \psi . \end{aligned} \quad (4.3)$$

These are slightly modified Coleman relations, or the ψ -equation and the ϕ -equation. The latter one is just the normal component of the axial current bosonization relation

$$\partial^\mu \phi = \sqrt{\pi} \bar{\psi} \gamma_\mu \gamma_5 \psi . \quad (4.4)$$

As known, in the two-dimensional world the axial current can be expressed easily through the vector current.

The tangential component of the bosonization relation is nicely presented in the “criminal story” of Nielsen and Wirzba⁴⁶. CCP is brought in while one is trying to derive the ψ -equation. In order not to get a decoupled ψ -equation, one should take care to allow the vanishing of the determinant of a 2×2 matrix, whose entries are derived by imposing locality, renormalizability and hermiticity conditions. For details see the work of Nadkarni and Nielsen⁴⁵. It happens that one finds solutions only for spacelike bag normals, i.e., timelike bag walls. This important constraint is automatically satisfied if the bag has AdS geometry, because of the timelike surface at spatial infinity ($\rho = \pi/2$) of AdS or CAdS. In their seminal paper Avis, Isham

and Storey have introduced reflective boundary conditions (RBC), later used extensively in AdS SUSY, but also “transparent” boundary conditions (TBC), which in our opinion are perfectly suited for discussing CCP. Imposing proper boundary conditions is essential for a good quantization scheme in AdS space. The difficulties with quantization in AdS background are related to the fundamental periodicities of AdS. As AdS is not globally hyperbolic, one has to pass to another conformally flat space, the Einstein static universe (ESU), proceed with quantization there, and only then mapping back to AdS. The detailed analysis ²⁶ shows that Cauchy data must be specified on two spacelike surfaces separated in time by π . The periodicities of the mode functions in AdS and ESU imply that only the following sum of “conserved” quantities is really constant

$$P_a = Q_a(\tau) + Q_a(\tau + \pi) , \quad (4.5)$$

where

$$Q_a(\tau) = \int_{\tau=\text{const}} T_{0\nu} \xi_a^\nu g^{00} \sqrt{-g} d\rho . \quad (4.6)$$

The symbols of energy-momentum tensor, Killing vectors and metric tensor are the standard ones.

The unusual conservation law (4.5) dictates the recirculation of the energy, angular momentum and conserved conformal quantities lost through the timelike interface at spatial infinity. In two dimensions the AdS background has SO(2,1) invariance and the conserved quantities correspond to the “translation” generator, dilatation, and restricted spatial conformal transformation, that is to the three conformal Killing vectors of the background geometry.

In curved space bosonization rules have to be modified in order to compensate the scale change of fermion masses by a position-dependent interaction with which the bosons have to be provided. Thus for massive fermions the ϕ -equation turns into

$$n\partial\phi = \sqrt{\pi} \Omega \bar{\psi} \not{n} \gamma_5 \psi , \quad (4.7)$$

where the conformal factor $\Omega = (\cos \rho)^{-1}$ for AdS.

A related problem is that of Dirac operators with space-dependent masses ⁴⁷.

It is worth noting that the helicity eigenstates in AdS are linear combinations of antipodal eigenstates ⁴⁸. The antipodal reflection is a spatial reflection plus a translation in time by $\Delta t = \pi/a$. For a more intuitive analogy, the antipodal reflection is similar to the total reflection in which the electromagnetic field is leaking in the propagation-“forbidden” medium causing a time “delay”. In this way one will not encounter with AdS bags a helicity paradox as for MIT bags (a boundary changes always the helicity) since “AdS-quarks” at the “walls” suffer antipodal reflections, a very appealing general fact.

5. The pion and the AdS bag

Theoretically, the pion has many faces ⁴⁹. In QCD, the pion may be considered as a single bound pair of valence quark and antiquark. Since u and d quarks are

almost massless on hadronic scales, the resulting chiral $SU(2)_L \times SU(2)_R$ symmetry of QCD suggests the identification of the pion with the associated Goldstone boson. It can be viewed also as a collective $q\bar{q}$ mode built on the QCD vacuum ⁴⁹.

The most attractive model remains that of Nambu & Jona-Lasinio (NJL). In this well-known model, the chiral symmetry is dynamically broken through an effective chiral interaction which moves the mass of the pionic $q\bar{q}$ mode down to zero according to the Goldstone theorem.

Many years ago, Castell has considered the problem of the Goldstone boson in AdS space ⁵⁰. He also introduced an invariant partial causal order possible only for AdS and not for dS. This time order is given for $s^2 \leq 0$ events by the Castell function

$$\mathcal{C}(x, x') = \text{signum}(x_0 x'_4 - x_4 x'_0) , \quad (5.1)$$

with the order values ± 1 for “before” and “after”. The Laplace-Beltrami equation has been written by Castell in AdS horospherical coordinates and he made use of a scheme of the Goldstone theorem worked out by Bludman and Guralnik. The result of Castell is that the Goldstone AdS boson is pseudoscalar and has a mass

$$m_{G,C} = m_\pi = 3a . \quad (5.2)$$

Castell’s AdS pion (the representation $D(3,0)$) is the Goldstone boson of the spontaneously broken conformal symmetry $SO(4,2) \rightarrow SO(3,2)$. The idea of connecting the pion to the breaking of conformal symmetry was reinforced by Isham, Salam and Strathdee ⁵¹ in an effective Lagrangian formalism in which the symmetry breaking term is taken to be in an irreducible representation of the combined conformal and chiral groups. In such a model the bare masses of the pion and dilaton field (introduced to make the Lagrangian conformal invariant) are connected.

Since in the Skyrme model the baryon charge has topological origin we have been extremely careful with the details of the Goldstone theorem. According to Shamir and Park ⁵², topological symmetries are never broken spontaneously whenever a local operator is used as the order parameter. Evading this by means of a delocalized operator, it can be shown that the “symmetry breaking” condition reflects the effect of the non-local order parameter on the boundary conditions at spatial infinity. A substantial part of the norm of the order parameter is to be found at spatial infinity.

The effects of a constant spacetime curvature on Nambu-Goldstone bosons were studied by Inami and Ooguri ⁵³, who showed that even in AdS_2 a continuous symmetry may be broken because the gravitational potential of this space, which confines particles within a finite spatial interval, is acting also as an infrared cutoff. The fluctuations of the Nambu-Goldstone mode decay exponentially near the spatial infinity. Also Verbin ⁵⁴ has dealt with the spontaneous symmetry breaking in AdS background of an $O(2)$ scalar doublet and obtained a massive Goldstone mode too.

6. Conclusions and perspectives

The ultimate goal of Skyrmion physics is the explanation of atomic nucleus

on a much more sophisticated basis than the conventional nuclear physics ⁵⁵, a commitment from the side of Mathematics.

One of the best candidates to achieve this difficult task is a hybrid skyrmion. However, in our opinion, it is mandatory that the bag inside the Skymion to be of AdS type and thus allow cosmology to enter the field. With the AdS bag the requirements of CCP are trivial and many other results are foreseeable. Surely the boundary of AdS bag, spreading all over an $S^1 \times S^2$ membrane, will be of great importance in the future Skyrmin physics. It is still to be decided if this membrane is leptodermous or pachidermous. It is well known that totally antisymmetric tensor fields as well as gravitational fields are able to produce closed membranes by quantum processes, such as instanton tunneling ⁵⁶. The interesting fact of the process is that it could stop automatically as soon as the effective cosmological constant becomes less or equal to zero. I believe this “neutralization of the cosmological constant by membrane creation” ⁵⁶ to be essential for further considerations on the lines of the present paper.

One may argue that because the spatial sections of AdS bubbles are infinite in extent these are not the right bags to put into the skyrmions. This objection is easy to reject. Heckman and Schücking have shown in 1959 that the space $\mathcal{H}^3$ could be substituted by a compact hyperbolic manifold iff some points are properly identified ⁵⁷. The resulting hyperbolic compact spaces have no continuous group of isometries as stated by a theorem of Yano and Böchner ⁵⁸. Killing vectors could be found at every point of these spaces but it is only a local property and no global Killing vector does exist. This is not an essential difficulty in femtocosmology. One should keep in mind also that in doing quantum field theory on such spaces confined by means of topological constraints, zero point energy phenomena (Casimir energy) are extremely important ⁵⁹.

One expects the bags to become more relevant in relativistic and ultrarelativistic nucleus-nucleus collisions, even though they are by far more appropriate for static conditions ⁶⁰. Some hints for the bags to be AdS spaces come from the hadronic multiplicity distributions where data are explained by parton branching probabilities ⁶¹. These probabilities have hypergeometric functions in their integral kernels and very similar hypergeometric functions enter the expressions for all sorts of field propagators in dS spaces ⁶². Moreover, a connection between QCD and femtocosmology is suggested by some works of Calzetta et al. ⁶³, in which they showed that the asymptotic freedom could be obtained by means of high curvature.

Even a torsion field may be accomodated. P. Minkowski ⁶⁴ has shown how little the Einstein equations are modified if one introduces a contorsion field of the form

$$K_{abc} = \epsilon_{abc} K \quad (a, b, c, = 1, 2, 3) , \quad (6.1)$$

where

$$K = \eta/R \quad (6.2)$$

with η an arbitrary constant and R the scale factor of the universe. The only effect of such a field is to change the curvature index parameter k ($k=1,0,-1$ for spherical,

flat and hyperbolic universes respectively) to the new value

$$k_1 = k - \eta^2 \quad (6.3)$$

A hyperbolic universe remains hyperbolic in the presence of a contorsion field of the type (6.3). This is of course an interesting result. Indeed, we could obtain hyperbolic femto-universes from a spherical universe by locally creating contorsion fields with the contorsion index $\eta = \pm\sqrt{2}$. Starting with a flat universe, we need the local contorsion index to be ± 1 .

Some other useful references are recorded at the end of the published ones ⁶⁵.

Acknowledgments

The author would like to thank Professor Abdus Salam, the International Atomic Energy Agency and UNESCO for hospitality at the International Centre for Theoretical Physics, Trieste.

References

1. A.W. Thomas, "The Quark Structure of Matter", PTP Suppl. **91**, 204 (1987)
 2. - V.I. Sanyuk, "Genesis and Evolution of the Skyrme Model from 1954 to the Present", preprint SU-4228-465 (1990)
 - E.H. Lieb, "Remarks on the Skyrme Model", Lecture given at the 1990 Summer Institute on Differential Geometry (1990)
 - C.L. Terng, "A Higher Dim. Generalization of the SG Eq. and its Soliton Theory", Ann. Math. **111**, 491 (1980)
 - N. Joshi, "An Explicit One-Soliton Solution of the 2+1 Dim. Generalized SG Eqs", Lett. Math. Phys. **20**, 261 (1990)
 - L.D. Fadeev, Lett. Math. Phys. **1**, 289 (1976)
 - Y. Yang, J. Math. Phys. **30**, 824 (1989)
 - I. Zahed, G.E. Brown, "The Skyrme Model", Phys. Rep. **142**, 1-102 (1986)
 - P.J. Mulders, Phys. Rev. **D30**, 1073 (1984)
 3. - E. Witten, Nucl. Phys. **B160**, 57 (1979)
 - L.-H. Chan, Phys. Rev. Lett. **55**, 21 (1985); **57**, 1199 (1986)
 4. - T.H.R. Skyrme, Proc. Roy. Soc. **A260**, 127 (1961); Nucl. Phys. **31**, 5 (1962)
 - H. Date, K. Fujii, and H. So, Lett. Math. Phys. **13**, 195 (1987)
 5. - E. Witten, Nucl. Phys. **B223**, 422 (1983)
 - S. Vadia, in *Lect. Notes in Phys.* **208** (Springer, 1984)
 6. M. Oka and A. Hosaka, "Nucleons as Skyrmions", to appear in The Annual Review of Nuclear and Particle Science, vol 42.
 7. G.E. Brown, in *Proc. Int. Conf. Nucl. Phys.*, Florence, Italy (1983)
 8. S. Nadkarni, H.B. Nielsen, and I. Zahed, Nucl. Phys. **B253**, 308 (1985)
 9. H. Rosu, preprint FT-337 (1988); Rev. Roum. Phys. **34**, 1239 (1988)
 10. - A.S. Eddington, *The Mathematical Theory of Relativity* (CUP, 1930)
 - R.C. Tolman, *Relativity, Thermodynamics and Cosmology* (OUP, 1934)
 - L. Godeaux, *Les Geometries Cayleyennes et Les Universes d'Einstein et de de Sitter* (Sciences et Lettres, 1947)
 - E. Schrödinger, *Expanding Universes* (CUP, 1956)
- Schrödinger discusses in his book the elliptic interpretation obtained by identifying the

- antipodal points. In group theory terms, one would say that the AdS rotations have imprimitivity classes containing pairs of antipodal points.
- V.A. Fock, *The Theory of Space, Time and Gravitation* (Pergamon Press, 1959)
 - H.P. Robertson and T.W. Noonan, *Relativity and Cosmology* (Saunders, 1968)
 - W. Rindler, *Essential Relativity* (Van Nostrand, 1969)
 - S. Weinberg, *Gravitation and Cosmology* (Wiley, 1972)
 - S.W. Hawking and G.F.R. Ellis, *The Large Scale Structure of Space and Time* (CUP, 1973)
 - N.D. Birell and P.C.W. Davies, *Quantum Fields in Curved Space* (CUP, 1982)
 - N.A. Gromov, *Contractions and Analytic Continuations of Classical Groups: A Unifying Treatment* (in Russian, Syktyvkar, 1990)
 - F. Gürsey, in *Group Theory Concepts and Methods in Elem. Part. Phys.*, (Gordon and Breach, 1964)
 - I.M. Yaglom et al., *Usp. Mat. Nauk* **19**:5, 51 (1964)
 - R.I. Pimenov, *Litov. Mat. Sb.* **5**:3, 457 (1965)
 - W. Thirring, *Acta Phys. Austr. Suppl. IV*, 269 (1967)
 - M. Gutzwiller, *Helv. Phys. Acta* **29**, 313 ((1956)
 - N.A. Chernikov and E.A. Tagirov, *Ann. Inst. H. Poincaré* **9**, 109 (1968)
 - O. Nachtmann, *Comm. Math. Phys.* **6**, 1 (1967)
 - G. Börner and H.P. Dürr, *Nuovo Cimento* **64A**, 669 (1967)
 - O. Phillips and E.P. Wigner, in *Group Theory and its Applications*, ed. E.M. Loeb (Academic Press, 1968)
 - B.S. De Witt, *Phys. Rep.* **19C**, 295 (1975)
 - E.W. Mielke, *Fort. d. Physik* **25**, 401 (1977)
 - G. Gensing, *J. Phys.* **A10**, 1687 (1977)
 - M.S. Drew, *Ann. Phys.* **103**, 469 (1977)
 - R. Drechsler and R. Sasaki, *Nuovo Cimento* **46 A**, 527 (1978)
 - M. Henneaux and C. Teitelboim, *Commun. Math. Phys.* **98**, 391 (1985)
 - F.J. Herranz, M.A. del Olmo and M. Santander, “Cayley-Klein Geometries Revisited”, presented at the II Int. Wigner Symp. (Goslar, 1991)
 - 11. P. Kerszberg, “The Einstein-deSitter Controversy of 1916-1917 and the Rise of Relativistic Cosmology”, in *Einstein Studies vol.1*, pp 325-366, Eds. D. Howard and J. Stachel (Birkhauser, 1989)
 - 12. J. Milnor, “Hyperbolic Geometry: The First 150 Years”, *Bull. AMS* **6**, 9 (1982)
 - 13. - J. Eiesland, *Trans. AMS*, **213**, 213 (1925)
 - S. Ianus, *St. Cerc. Mat.* **17**, 1161 (1965)
 - 14. B. Binetgar, C. Fronsdal and W. Heidenreich, *Ann. Phys.* **149**, 254 (1983)
 - 15. G.W. Gibbons, Aspects of Supergravity Theories - Three Lectures in Spain, in *Supersymmetry, Supergravity and Related Topics*, eds. F. del Aguila, J.A. de Azcárraga and L.E. Ibáñez (World Scientific, Singapore, (1985), p. 123
 - 16. W. Heidenreich, *Phys. Lett.* **B110**, 461 (1982)
 - 17. M.J. Duff, B.E.N. Nilsson, C.N. Pope, *Phys. Rep.* **130**, 1 (1986)
 - 18. E. Angelopoulos, in *Quantum Theory, Groups, Fields and Particles*, Ed. A.O. Barut, pp. 101-148 (Reidel, 1983)
 - 19. H. Nicolai, preprint CERN TH-3882 (1984)
 - 20. P.A.M. Dirac, *J. Math. Phys.* **4**, 901 (1963)
 - 21. - C. Fronsdal, *Phys. Rev.* **D26**, 1988 (1982); **D35**, 1262 (1987)
 - M. Flato and C. Fronsdal, *J. Math. Phys.* **22**, 1100 (1981)
 - 22. M.J. Duff, “Supermembranes: The First 15 Weeks”, CERN-TH-4797 (1987)
 - 23. E. Bergshoeff et al., *Phys. Lett.* **B199**, 69 (1987)
 - 24. - M. Flato and C. Fronsdal, *Phys. Lett.* **B172**, 412 (1986); *Lett. Math. Phys.* **20**, 65

- (1990); J. Math. Phys. **32**, 524 (1991)
 - M. Nakashima, Lett. Math. Phys. **18**, 237 (1988)
25. - L. Castell and W. Heidenreich, Phys. Rev. **D24**, 371 (1981); Acta Phys. Austriaca Suppl. XXXIII, 605 (1981)
 - H. Rumpf, Phys. Rev. **D24**, 275 (1981)
26. S.J. Avis, C.J. Isham, and D. Storey, Phys. Rev. **D18**, 3565 (1978)
27. A short list of papers is the following:
 - C.J.C. Burges et al., Ann. Phys. **167**, 285 (1986)
 - B. Allen et al., Phys. Lett. **B189**, 304 (1987)
 - D. Bernard, A. Folacci, Phys. Rev. D34, 2286 (1986)
 - R. Floreanini, C.T. Hill and R. Jackiw, Ann. Phys. **175**, 354 (1987)
 - P. Breitenlohner and D.Z. Freedman, Ann. Phys. **144**, 249 (1982)
 - S. Deser and R. Nepomechie, Ann. Phys. **154**, 396 (1984)
 - T. Inami and H. Ooguri, Progr. Theor. Phys. **73**, 1051 (1985)
 - E.S. Fradkin and M.A. Vasiliev, Ann. Phys. **177**, 63 (1987)
28. A. Bohm, Phys. Rev. **D33**, 3358 (1986)
29. K.S. Stelle and P.C. West, Phys. Rev. **D21**, 1466 (1980)
30. W. Miller Jr., *Symmetry and Separation of Variables*, Enciclop. of Math. and its Appl., Ed. Gian-Carlo Rota (Addison-Wesley, 1977)
31. B. Allen and M. Turyan, Nucl. Phys. B292, 813 (1987)
32. - C. Sivaram and K.P. Sinha, Phys. Rep. **51**, 111 (1979)
 - S.I. Fisenko et al., Phys. Lett. **A148**, 405 (1990)
 - E. Recami and V. Tonin-Zanchin, preprint INFN/AE - 91/103 (1991)
 - M. Kovalyov, Hadronic J. **12**, 51 (1989)
33. A. Salam and J. Strathdee, Phys. Rev. **D18**, 4596 (1978)
34. R. Prasad, Nuovo Cimento **44A**, 4327 (1966)
35. - Z. Haba, Phys. Lett. **B78**, 421 (1978)
 - C.C. Bernido, preprint IC/85/75 (1985)
36. M. Romaniuk, in Proc. VIII Warsaw Symp. on Elem. Part. Phys., Kazimierz, Poland, p. 263 (1985)
37. E. Van Beveren, C. Dullemond and T.A. Rijken, Phys. Rev. **D30**, 1103 (1984)
38. -N. Rosen, Gen. Rel. Grav. **12**, 493 (1980)
 - C. Dullemond, Gen. Rel. Grav. **20**, 139 (1988)
39. H.J. Bhabha, "The Fundamental Length introduced by the Theory of the Mesotron", Nature (Suppl), p. 276 (18 Febr. 1939)
40. S. Fubini, Nuovo Cimento **34A**, 521 (1976)
41. P. Roman and J. Haavisto, Int. J. Theor. Phys. **16**, 915 (1977)
42. S. Coleman, Phys. Rev. **D11**, 2088 (1975)
43. V.A. Rubakov, Nucl. Phys. **B236**, 109 (1984)
44. - M. Rho, Preprints SPhT/86-159 (1986), SPhT/84-123 (1984)
 - R.J. Perry and M. Rho, Phys. Rev. **D34**, 1169 (1986)
45. - S. Nadkarni and H.B. Nielsen, Nucl. Phys. **B263**, 1 (1986)
 See also the following references:
 - B.-Y. Park, V. Vento, "Proton Spin and the CCP", Nucl. Phys. **A513**, 413 (1990)
 - H.B. Nielsen et al., "Color Anomaly in a Hybrid Bag Model", Nordita-91/4 NP (1991)
 - *Workshop on Skyrmions and Anomalies*, Krakow, (World Scientific, 1987)
46. H.B. Nielsen and A. Wirzba, preprint NBI-HE-87-32 (1987)
47. J.Z. Imbrie et al., Commun. Math. Phys. **135**, 421 (1991)
48. W. Heidenreich, Phys. Rev. **D36**, 1685 (1987)
49. - W. Weise, Nucl. Phys. **A434**, 685c (1985)
 -A.W. Thomas, Nucl. Phys. **A518**, 186 (1990)

50. L. Castell, Nuovo Cimento, **61A**, 585 (1968)
See also:
G. Börner, Prog. Theor. Phys. **43**, 244 (1970)
51. C.J. Isham, A. Salam, and J. Strathdee, Phys. Rev. **D2**, 685 (1970)
52. Y. Shamir and S.H. Park, Phys. Lett. **B258**, 179 (1991)
53. T. Inami and H. Ooguri, preprint RIFP-612 (1985)
54. Y. Verbin, Nucl. Phys. **B272**, 739 (1986); Phys. Lett. **B197**, 55 (1987)
55. I. Klebanov, Nucl. Phys. **B262**, 133 (1985)
56. - J. David Brown and C. Teitelboim, Nucl. Phys. **B297**, 787 (1988)
- Yu. E. Pokrovskii, Yad. Fiz. **50**, 907 (1989)
- A.T. Barnaveli, O.V. Kancheli, Sov. J. Nucl. Phys. **52**, ?
- H. Sato, "Motion of a Shell at a Metric Junction", PTP **76**, 1250 (1986)
- J. Ipser and P. Sikivie, "Gravitationally Repulsive DW", Phys. Rev. **D30**, 712 (1984) 494 (1990)
57. - O. Heckman and E. Schücking, in Hand. d. Physik **53**, p. 515 (Springer, 1959)
See also the following list of mathematical papers:
- H.V. Fagundes, "Closed Spaces in Cosmology", preprint IFT-P.013/91 (1991); Lett. Math. Phys. **6**, 417 (1982)
- G.F.R. Ellis, Gen. Rel. Grav. **2**, 7 (1971)
- L.A. Best, Can. Journ. Math. **23**, (3), 451 (1971)
- E.M. Andreev, Matem. Sb. **81**, 445 (1970)
- E.B. Vinberg, Usp. Mat. Nauk **40**, 29 (1985)
- I.S. Gutsul, Trudy Mat. Inst. V.A. Steklov **152**, 89 (1980)
- A.Yu. Vesnin, Sib. Mat. J., **28**:5, 50 (1987)
- T. Jorgensen, Ann. Math. **106**, 61 (1977)
- C. Weber, H. Seifert, "Die beiden Dodekaederräume", Math. Zf. **37**, 237 (1933)
- P. Scott, "The geometries of 3-manifolds", Bull. Lond. Math. Soc. **15**, 401 (1983)
- C. Series, "Non-Euclidean Kaleidoscopes", Interdisciplinary Science Reviews **15**, 224 (1990)
- V.P. Frolov, Phys. Rev. **D43**, 3878 (1991)
In Frolov's paper the membrane $S^1 \times S^2$ is treated as a compactified 3d space with one wormhole, being an example of a time machine.
- G. D. Mostow, Bull. Am. Math. Soc. **16**, 225 (1987)
- J.P. Gaspard, R. Mosseri, and J.F. Sadoc, Phil. Mag. **B45**, 467 (1982); R. Mosseri and J.F. Sadoc, Phil. Mag. **B50**, 557 (1984)
- A.N. Cyuvyrov et al., JETP **89**, 2052 (1985)
- I. A. Ovidko, Ukr. Fiz. J. **34**, 1343 (1989)
- A.M. Srivastava, Phys Rev. **D43**, 1047 (1991)
58. K. Yano and S. Böchner, *Curvature and Betti Numbers* (PUP, 1953)
59. R.T. Hammond and W.A. McKinley, Phys. Rev. **D20**, 3047 (1979)
60. - V.G. Kadyshevsky and D.V. Fursaev, Teor. Mat. Fiz. **83**, 197 (1990)
- G.A. Gal'perin, Dokl. Ak. Nauk **302**, ? (1988); [Sov. Math. Dokl. **38**, 367 (1989)]
- Yu.A. Golfand, JETP **43**, 256 (1962)
61. B. Durand and I. Sarcevic, Phys. Rev. **D36**, 2693 (1987)
62. B. Allen and T. Jacobson, Comm. Math. Phys. **103**, 669 (1986)
63. E. Calzetta, I. Jack and L. Parker, Phys. Rev. Lett. **55**, 1241 (1985)
64. P. Minkowski, Phys. Lett. **B173**, 247 (1986)
65. **More References - added in 1994**
R. Shankar, "Bosonization: How to make it work for you in condensed matter", Lectures at BCSPIN School, Katmandu, May 1991
C. Bonatti, "Un point fixe commun pour des diffeos commutants de S^2 ", Ann. Math.

129, 61 (1989) and refs therein.
 B. Konopelchenko and C. Rogers, “On (2+1)-dim nonlinear systems of Loevner type”, Phys. Lett. **A158**, 391 (1991)
 J.D. North, *The Measure of the Universe, A History of Modern Cosmology*, (Dover 1990, first published in 1965 by OUP)
 This very good book has four paragraphs on de Sitter worlds as follows: The de Sitter world and Mach’s Principle, The de Sitter effect and Weyl’s Principle, Material Content, The geometry of the de Sitter world.
 Federico Zertuche Mones, “2+1 Quantum de Sitter Gravity”, Ph.D. Thesis at ISAS-Trieste (Academic Year 1988-1989)
 M. Maischak, “The Gravitational Anomalies in de Sitter Space”, ITP-UH 8/91, Hannover (1991)
 H.-J. Schmidt, “On the de Sitter space-time: the geometric foundation of inflationary cosmology”, PRE-ZIAP 91-04, Fortschr. Phys. (1992)
 M. Chaichian et al., “Oscillator realization of the q-deformed anti-de Sitter algebra”, Phys. Lett. **B291**, 411 (1992)
 J. Ellis et al., “Quark solitons as constituents of hadrons”, CERN-TH-6426/92 (1992)
 H.S. Sharatchandra, “Bosonisation in any dimension”, Preprint No. imsc-92 (Madras)
 V.K. Dobrev and P.J. Moylan, “Finite-dimensional singletons of the quantum anti de Sitter algebra”, Phys. Lett. **B315**, 292 (1993)
 I.G. Moss and S.J. Poletti, “Conformal anomalies on Einstein spaces with boundary”, NCL94-TP10 (1994)
 S. Alexander Ridgway, “ $S^1 \times S^2$ wormholes and topological charge”, Phys. Rev. **D50**, 892 (1994)
 D. Giulini, “3-manifolds for relativists”, Int. J. Theor. Phys. **33**, 913 (1994)
 S. Carlip and S.P. de Alwis, “Wormholes in 2+1 dimensions”, Nucl. Phys. **B337**, 681 (1990)
 N.S. Manton, Commun. Math. Phys. **111**, 469 (1987)
 A. Wirzba, “Skyrmions on the hypersphere”, preprint Nordita.